Using the Selective Functional Movement Assessment for the Evaluation of Dancers' Functional Limitations and Dysfunctions: A Critically Appraised Topic

Victoria Fauntroy, Marcie Fyock, Jena Hansen-Honeycutt, Esther Nolton,
and Jatin P. Ambegaonkar

Clinical Scenario: Dancers participate in a functionally demanding activity. Athletic participation typically requires the completion of a preparticipation examination, which involves a functional movement screen offering insight into potential injury recognition. The Selective Functional Movement Assessment (SFMA) was created to measure the status of movement–pattern-related pain and dysfunction using regionally interdependent movement to aggravate symptoms and exhibit limitations and dysfunctions. Still, a functional assessment has not been identified to recognize potential dysfunctions or limitations in this population. **Clinical Question:** Does the use of the SFMA improve overall evaluation of dancers by providing more information on a dancer's overall functional ability and limitations? **Summary of Key Findings:** The literature search discovered 12 studies and 3 books in which 4 studies were included (2 case reviews, 1 case report, and 1 original research study) based on the inclusion and exclusion criteria. Three of the studies provided clinical case studies utilizing the SFMA to improve the patient's dysfunctions, whereas 1 study examined the intrarater and interrater reliability of the SFMA. In 3 studies, participants displayed less movement dysfunction. The authors from 3 of the studies agreed the SFMA was a valuable tool for clinicians to use during evaluations, as it provided a more holistic view of the patient, discovering dysfunctional movement patterns that may better identify the source of injury. **Clinical Bottom Line:** Low-quality evidence, defined as poorly designed case studies, case series, and cohort studies, exist that supports improvement of overall evaluations when utilizing the SFMA. Although the studies were considered low-quality evidence, each included study displayed an effective use of the SFMA as an overall evaluation that correctly identified dysfunctional movement patterns. **Strength of Recommendation:** Grade C evidence exists that the SFMA contributes to the functional evaluation used in dancers.

Keywords: SFMA, athlete, regional interdependence, preparticipation examination

Clinical Scenario

Historically, athletic participation required the completion of a functional movement screening during the preparticipation examination, which provided valuable information to clinicians regarding injury prevention. However, in recent years, functional movement screenings have begun to be incorporated in the preparticipation examination, providing additional injury recognition information. Recently, the Selective Functional Movement Assessment (SFMA) was developed by a team of clinicians and has its origins in the functional movement screen (FMS) to measure the status of movement–pattern-related pain and dysfunction.^{1,2} The SFMA uses regionally interdependent movement to exacerbate symptoms leading to the manifestation of limitations and dysfunctions, with the resulting information relating to the patient's chief complaint in the attempt to create more functional movement and decrease injury.^{2,3}

Traditionally, clinicians have utilized a biomedical model of disease as the basis of evaluation, focusing solely on a biological reason to explain injury or illness.⁴ The biomedical model often focuses treatment at the location of pain or chief complaint and

rarely considers other anatomical locations as possible reasons for the injury. The limited information provided by this model narrowed the available treatment options. Recently, a paradigm shift to expand on the biomedical model and include other factors or regions that may contribute to the individuals' complaint has occurred.⁴

The regional interdependence (RI) model describes the concept that a patient's primary musculoskeletal symptom(s) may be directly or indirectly related or influenced by impairments from various body regions and systems regardless of proximity to the primary symptom(s).⁴ The RI model presents an opportunity for clinicians to identify remote dysfunctions that may be affecting or causing the patient's chief complaint, which is an integral role in the SFMA. The SFMA is a clinical assessment that is designed to systematically identify causes of movement dysfunction while taking pain into consideration using an algorithmic approach.¹ Guided by the RI model, the SFMA can be used by the clinician to identify dysfunctions in different parts of the body and determine their relationship in the overall injury.

The SFMA has been used to identify remote dysfunctions through the utilization of 7 top-tier (ie, basic movement) assessments: cervical patterns—flexion, extension, and rotation (right [R] + left [L]); upper-extremity patterns: medial rotation–extension pattern (R + L) and lateral rotation–abduction pattern (R + L); multisegmental patterns: multisegmental flexion; multisegmental extension; multisegmental rotation (R + L); single-leg stance (R + L); and overhead squat.⁵ The results of the SFMA are separated into 4 categories: functional nonpainful, functional painful,

Fauntroy, Fyock, Nolton, and Ambegaonkar are with the Sports Medicine Assessment Research and Testing (SMART) Laboratory, George Mason University, Manassas, VA, USA. Hansen-Honeycutt is with the Department of Dance, George Mason University, Fairfax, VA, USA. Ambegaonkar (jambegao@gmu.edu) is corresponding author.

dysfunctional nonpainful, and dysfunctional painful.² The SFMA has also been used to evaluate athletes in various sports such as soccer, weight lifting, running, and baseball.⁵⁻⁷

However, a functional assessment has not been identified for dancers. Dancers participate in a physically demanding activity requiring extreme functional movement. As with any sport, dancers suffer injuries resulting from the demands required by their activity and will often continue to train while in pain. Specifically, dancers have high rates of injury, with 80% to 97% of university, professional ballet, and modern dancers reporting at least 1 musculoskeletal injury per year.⁸ Most of these injuries occur in the lower-extremity or lumbar spine because of overuse or repetitive motions.⁸⁻¹⁰ Thus, there is a necessity for the critical appraisal of research to examine the effectiveness of using the SFMA in the evaluation process to assess dancers' functional movement to decrease injury incidence. If the true dysfunction causing the injury is not addressed, other regions of the body will compensate to produce the movement required for the activity.⁵ Compensation may only lead to further injuries and pain for the dancer.

Focused Clinical Question

Does the use of the SFMA improve overall evaluation of dancers by providing more information on the dancers' overall functional ability and limitations?

Summary of Search, "Best Evidence" Appraised, and Key Findings

- Twelve studies and 3 books were identified during the electronic search; 2 case reviews, 1 case report, and 1 original research study fulfilled the inclusion and exclusion criteria and were included in this critically appraised topic (CAT) (Table 1).
- Three of the included studies provided clinical case studies using the SFMA with appropriate treatment interventions to improve the patient's dysfunctions.^{5,6,10}
- One included study examined the intrarater and interrater reliability of the SFMA using both the categorical scoring tool and criterion checklist in raters with various levels of experience.¹¹
- In 3 of the studies included in this CAT, all dysfunctional movement demonstrated by the participants was resolved and completed in a more functional manner.^{5,6,10}
- In the last included study, the authors found that the intrarater and interrater reliability of categorical scoring and criterion checklist scoring of the SFMA were reflective of level of experience. Specifically, in a healthy population, the SFMA was most reliable between sessions when performed by a single-experienced rater.¹¹
- The authors from the included studies that used the SFMA agreed that the SFMA was a beneficial tool for clinicians during evaluations because it provided a more integrated view of the patient, discovering dysfunctions that may have been the cause of injury.^{5,6,10,11}

Clinical Bottom Line

There is low-quality evidence for the improvement of overall evaluations using the SFMA based on the level 4 evidence of

Figure 1 — Summary of search strategy.

the studies reviewed for this CAT. Although the studies were considered low-quality evidence, each study concluded that the SFMA correctly identified dysfunctions and limitations causing the injury, which assisted the clinicians in the creation of proper rehabilitation programs for their patients. Nevertheless, because of limited original research and/or high-quality studies that discuss the SFMA only, this CAT reviewed studies deemed poor-quality evidence to answer the clinical question.

Strength of Recommendation

Grade C evidence exists that the SFMA improves the overall evaluation of dancers by contributing additional information on the comprehensive functional ability and limitations of the dancer. The Oxford Centre for Evidence-Based Medicine defines level 4 evidence as poorly designed case studies, case series, and cohort studies. A grade of C is suggested by the Oxford Centre for Evidence-Based Medicine for level 4 studies.¹²

Search Strategy

An electronic search was conducted in September and October 2017 (Figure 1).

Terms Used to Guide Search Strategy

- Patient/Client Group: All dancers (ie, no specific discipline)
- Intervention (or Assessment): Selective Functional Movement Assessment (SFMA)

- **Comparison:** Athletes participating in competitive sport (ie, team sports)
- **Outcome(s):** Improved overall evaluation

Sources of Evidence Searched (Databases)

- CINAHL Plus
- MEDLINE
- PubMed Central
- Primo by Ex Libris
- Supplementary resources retrieved via reference list review and web searches

Inclusion and Exclusion Criteria

Inclusion Criteria

- Restricted to studies describing SFMA and its uses in dancers
- Restricted to studies in the English language
- Restricted to studies published within in the last 7 years (2010–2017)

Exclusion Criteria

- Systematic reviews, meta-analyses, clinical reviews, books, CATs, and clinical commentaries

Results of Search

Four pertinent studies were discovered and classified as presented in Table 1. The validity of the selected studies was assessed using the Physiotherapy Evidence Database (PEDro) scale to score each study based on certain criteria. The included studies also received a letter grade and level of evidence number as defined by the Oxford Centre for Evidence-Based Medicine.¹²

Best Evidence

The ensuing studies were recognized as the “best” evidence and chosen for inclusion in the CAT. The reasons for selecting these studies were as follows:

- Met all inclusion and exclusion criteria
- Discussed SFMA as an evaluation tool

Table 1 Summary of Study Designs of Articles Using the Selective Functional Movement Assessment to Measure the Status of Movement Pattern-Related Pain and Dysfunction

Level of evidence	Study design/ methodology of articles retrieved	Number located	Author (year)
4	Case report	1	Goshtigian and Swanson (2016) ⁵
4	Case review	2	Krzyzanowicz et al (2015) ¹⁰ Mokha et al (2015) ⁶
3b	Original research	1	Glaws et al (2014) ¹¹

Summary of Best Evidence

Table 2 describes characteristics of included studies using the SFMA to measure the status of movement pattern-related pain and dysfunction.

Implications for Practice, Education, and Future Research

Dance is a physically demanding, athletic activity requiring elevated functional ability to repeatedly perform at a prominent level. For example, professional dancers routinely participate in technique classes during most of any given day, followed by rehearsals and performances through the late afternoons and evenings. Preparticipation exams are frequently performed in athletic sport participation, yet this is not commonly performed for dancers. A functional screening for dancers may be of benefit to observe the movement patterns of the dancer.¹⁰ Krzyzanowicz et al¹⁰ noted that the SFMA correctly identified dysfunctions leading to sacroiliac joint pain in 3 dancers. The authors suggested the use of the SFMA to other clinicians as part of their evaluation process and to help with the proper creation of rehabilitation programs to correct the dysfunction.

The SFMA incorporates the “breakout,” which is utilizing a movement pattern isolation map to establish probable causes for the dysfunction when a movement is not categorized as functional nonpainful.¹⁰ The dysfunction is divided into 3 categories: (1) tissue extensibility dysfunction (TED), (2) joint mobility dysfunction (JMD), or (3) stability or motor control dysfunction (SMCD).¹⁰ A tissue extensibility dysfunction recognizes tissues that cross more than 1 joint or are multiarticular (ie, neural tension, fascial tension, or muscle shortening).^{2,10} A joint mobility dysfunction classifies the articular surfaces and the contractile and noncontractile connecting spinal articular segments that exhibit decreased mobility (ie, osteoarthritis, adhesive capsulitis, fusion).² A stability or motor control dysfunction is broken into either a stability dysfunction (ie, strength) or a motor control dysfunction (ie, neurological processing problem).^{2,10}

Once the dysfunctions are recognized utilizing the SFMA breakout, therapeutic rehabilitation programs can be implemented that improve the dancer’s dysfunctional nonpainful or dysfunctional painful movement to functional nonpainful patterns. Interestingly, Krzyzanowicz et al¹⁰ noticed that each dancer exhibited dysfunction in the multisegmental flexion, single-leg stance, and overhead squat patterns. Goshtigian and Swanson⁵ discovered their college soccer player displayed dysfunction in the cervical rotation (R + L), medial rotation–extension, overhead squat, and all multisegmental (flexion, extension, rotation [R + L]) patterns. Similarly, Mokha et al⁶ recorded their college runner’s demonstrated dysfunction in all multisegmental patterns (flexion, extension, rotation [R + L]). In these cases, the discovery of these dysfunctions contributed information that helped decrease the patient’s chief complaint. The treatment administered to each patient was categorized based on the SFMA breakout findings. For example, patient 1 in the Krzyzanowicz et al¹⁰ study completed the rolling technique and other similar exercises to address her stability or motor control dysfunction,¹⁰ whereas the patient in the Goshtigian and Swanson⁵ study performed joint mobilizations to address his joint mobility dysfunction. The included case reviews and single case study provided vital information in this CAT that may allow for a more appropriate rehabilitation and treatment protocol. In the reviewed studies, all concluded the SFMA was beneficial and

Table 2 Characteristics of Included Studies Using the Selective Functional Movement Assessment to Measure the Status of Movement Pattern-Related Pain and Dysfunction

	Goshtigian and Swanson (2016)⁵	Krzyzanowicz et al (2015)¹⁰	Mokha et al (2015)⁶	Glaws et al (2014)¹¹
Study design	Case report	Case review	Case review	Original research
Participants	18-y-old male. DDx: facet joint dysfunction, intervertebral disk pathology, lumbar strain, SI joint dysfunction, hip muscular strain, impingement, and/or hip bursitis. Dx: DN in MSF, MSE, and MSR; cervical and UE, overhead deep squat. Hip TED, thoracic JMD, and SMCD.	Sample of convenience. 3 female dancers • Pt. 1: 18 y old. Dx: L innominate; ANT rotation. DN: R cervical rotation, MSF, MSR (R), single-leg stance (bilat.), and overhead squat. SMCD. • Pt. 2: 19 y old. Dx: L innominate; POST rotation. DP: MSF, MSE, and overhead squat. DN: MSR (R) and single-leg stance (bilat.). Extension JMD and SMCD. • Pt. 3: 18 y old. Dx: R innominate; ANT rotation. FP: cervical flexion and MSR (R). DN: MSF. DP: single-leg stance (R) and overhead squat. Hip JMD.	19-y-old female Diagnosis made by both an ATC and board certified orthopedic PT. Dx: ESHS; chronic R hip p; weak R hip ABD, ER, and extension; asymmetries in FMS rotary stability and inline lunge tests; incomplete patterns in the FMS; trunk stability push-up. DN in MSF, MSE, and MSR from the SFMA and SMCD. Running mechanics: hip ADD and IR, CPD, and rearfoot EVER. DDx: greater trochanteric bursitis, iliopsoas tendonitis, and intraarticular lesions.	Convenience sampling consisting primarily of PT students and OSU Club Rugby team. 39 healthy participants (27 males, 12 females). 35 participants (24 males, 11 females) involved in final statistical analyses resulting from data error in 4 subjects. Potential participants were excluded if they had undergone orthopedic surgery within the past 6 mo, were currently pregnant, recorded any positive marks on the PAR-Q health assessment, or were under the age of 18 y. 3 raters with varying levels of experience participated.
Intervention investigated	Subject given HEP Seen for 13 visits over 9 wk, ranging from 45 min to 1 h. Began with 1–2 visits per week, then 1 visit per week during last 3 wk. Interventions (based on Guide to Physical Therapist Practice): manual therapy, motor function training, and therapeutic exercises. PTs provided Tx. At beginning of every Tx session, subject assessed using each of SFMA top-tier movements that were DN during previous visit.	Mulligan MWM dependent on innominate rotation. Each performed 3 sets × 10 reps. Exercises selected were based on SFMA findings for each pt and were performed before the Mulligan MWM. Only DN patterns were reassessed at discharge. All patients were monitored for the rest of the academic semester. Doctoral AT student (with 7-y experience and 1 y using Mulligan concept) gave initial evaluation and subsequent exercises.	Attended 12 corrective exercise sessions over 4 wk with the AT or PT, lasting ~30 min each. Focus of all sessions was correcting functional movement pattern impairments as identified using SFMA. No corrective exercises directly targeted running technique.	Video recording: each participant during 7 top-tier assessments (10 movements). Raters not present during data collection and later individually evaluated each video using 2 scoring sheets: (1) categorical scoring sheet (from SFMA developers) and (2) 34-point criterion checklist scoring tool (SFMA manual). Evaluators repeated scoring process for intrarater reliability between 7 and 14 d following initial assessment. Test administrator: PT (8 h of didactic SFMA training) provided demonstration of each movement prior to completion of task. Evaluators' composite results compared within and between raters using absolute agreement and kappa coefficient based on categorical classification of each movement.
Outcome measure(s)	<i>Primary outcomes:</i> SFMA results <i>Secondary outcomes:</i> VAS, special test results, thoracic spine joint mobility	<i>Primary outcomes:</i> SFMA results, DPA scale scores, NPRS scores, special test results <i>Secondary outcomes:</i> DPA scale scores, NPRS scores, special test results.	<i>Primary outcomes:</i> SFMA and FMS results <i>Secondary outcomes:</i> VAS, MMT, special tests, running mechanics	<i>Primary outcomes:</i> Intrarater and interrater reliability <i>Secondary outcomes:</i> SFMA scores using categorical scoring sheet and 34-point criterion checklist

(continued)

Table 2 (continued)

	Goshtigian and Swanson (2016) ⁵	Krzyzanowicz et al (2015) ¹⁰	Mokha et al (2015) ⁶	Glaws et al (2014) ¹¹
Main findings	SFMA: 9/10 DN patterns to 1/10 DN Special tests: 3 pos. → neg. Hip strength: 4/5 (L) and 4+/5 (R) all planes → 5/5 (bilat.) Thoracic spine joint mobility: 2/6 → 3/6 Pain: 2–7/10 → 0–2/10	Pt. 1: • SFMA: 6/10 DN → FN • DPA: 31 to 2 • NRPS: 3/10 (dancing and rest) to 0/10 • Special tests: 2 pos.-SIJ pl; 1 pos.-ANT innominate rotation → neg. Pt. 2: • SFMA: 3/10 DP; 3/10 DN → FN • DPA: 49 to 4 • NRPS: 7/10 (dancing) to 0/10 • Special tests: 2 pos.-SIJ pl; 1 pos.-POST innominate rotation → neg. Pt. 3: • SFMA: 2/10 FP; 1/10 DN; 2/10 DP → FN • DPA: 15 to 9 • NRPS: 8/10 (dancing) to 1/10 • Special tests: 2 pos.-SIJ pl; 1 pos.-ANT innominate rotation → neg.	SFMA: 3/10 DN → FN FMS: 15/21 → 17/21 Pain: 4/10 → 0/10 Running mechanics: • Hip ADD: 19.3 (1.13) → 17.4 (0.21) (R); 16.3 (2.33) → 19.4 (1.13) (L) • Rearfoot EVER: 7.2 (0.0) → 3.7 (0.0) (R); 6.5 (0.99) → 6.9 (1.99) (L) • 76.5 N (15% BW) decrease in vertical ground reaction forces. • No clinically meaningful changes in R hip IR or CPD • Hip IR: 34.0 (1.48) → 19.4 (1.13) (L); 23.0 (0.49) → 23.2 (0.35) (R)	Intrater and interrater reliability of categorical scoring and criterion checklist scoring of the fundamental movements of the SFMA were reflective of level of experience. Fundamental SFMA movement scoring demonstrated substantial to excellent intrater reliability when using the categorical scoring tool but poor to good intrater reliability when using criterion-based scoring tool. Substantial and moderate interrater reliability was found for more experienced users with categorical and criterion scoring, respectively, whereas poor reliability was found for least experienced rater.
Level of evidence	4	4	4	3b
Validity score (if applicable)	5/10	5/10	5/10	4/10
Conclusion	The SFMA is effective as a method or system used to qualitatively analyze movement at initial examination and through the treatment process and to direct subsequent intervention choices. The SFMA helped therapists recognize dysfunctional movements that were present in subsequent regions that were not seen with more conventional examination procedures.	For this case series, the use of SFMA and Mulligan MWM interventions was associated with clinically significant improvements in pain using the NPRS, disability using the DPA scale, and functional movement using the SFMA for each patient. Given that 3 consecutive patients with SIJ pain of varying duration (3 d–12 mo) improved in an average of 4.3 treatments, it would suggest that this intervention strategy was successful.	Correcting faulty fundamental movement patterns had positive effects on reducing known running pathomechanics and relieving pain in this female competitive runner with ESHS.	Scoring of the 10 fundamental movements of SFMA, using the categorical and criterion checklist model in a healthy population, is most reliable between sessions when performed by a single-experienced rater.

Abbreviations: ABD, abduction; ADD, adduction; ANT, anterior; AT, athletic trainer; ATC, certified athletic trainer; bilat., bilateral; BW, body weight; CPD, contralateral pelvis drop; DDx, differential diagnosis; DN, dysfunctional nonpainful; DP, dysfunctional painful; DPA, disablement in the physically active; Dx, diagnosis; ER, external rotation; ESHS, external snapping hip syndrome; EVER, eversion; FMS, functional movement screen; FN, functional nonpainful; FP, functional painful; HEP, home exercise program; IR, internal rotation; JMD, joint mobility dysfunction; L, left; MMT, manual muscle test; MSE, multisegmental extension; MSF, multisegmental flexion; MSR, multisegmental rotation; MWM, Mulligan mobilizations with movement; neg., negative; NPRS, Numeric Pain Rating Scale; OSU, Ohio State University; pl, pain; PAR-Q, Physical Activity Readiness Questionnaire; pos., positive; POST, posterior; pt., patient; PT, physical therapist; R, right; reps., repetitions; SFMA, Selective Functional Movement Assessment; SI, sacroiliac joint; SMCD, stability or motor control dysfunction; TED, tissue extensibility dysfunction; Tx, treatment; UE, upper extremity; VAS, visual analog scale.

relevant to identifying both dance and other team sport athletes' movement dysfunction^{5,6,10,11} because the method incorporates the RI model to identify and treat dysfunctions in the body to provide holistic functional movement.⁴

Based on the conclusions of the authors in each of the included studies, there is moderate evidence to suggest that clinicians use the SFMA to improve a dancer's overall clinical evaluation. Although the studies were considered low-quality evidence, each included study displayed effective use of the SFMA as an overall evaluation that correctly identified dysfunctional movement patterns. Additional research needs to be conducted on the efficacy of the SFMA and its associated treatment outcomes, as well as the validity of the SFMA itself. Future studies should specifically concentrate on the use of the SFMA in the dance populations, as there is little published information about a dance-specific functional screening. Finally, this CAT should be reviewed in 2 years to determine whether additional best research evidence has been published that could aid in answering the focused clinical question about examining dancers' functional capacities with the goals of decreasing their injury risk and improving their performance.

Acknowledgments

The authors declare no conflict of interest.

References

1. Cook G, Burton L, Hoogenboom BJ, Voight M. Functional movement screening: the use of fundamental movements as an assessment of function—part 1. *Int J Sports Phys Ther*. 2014;9(3):396–409. PubMed ID: [24944860](#)
2. Cook G. *Movement: Functional Movement Systems: Screening, Assessment and Corrective Strategies*. Aptos, CA: Target Publications; 2010.
3. Rhinehart AJ, Schroeder KM, May J, Baker R, Nasypany A, Ambegaonkar JP. Movement assessment: techniques and possible integration into clinical practice. *Int J Athl Ther Train*. 2015;20(6): 5–9. doi:[10.1123/ijatt.2015-0021](#)
4. Sueki DG, Cleland JA, Wainner RS. A regional interdependence model of musculoskeletal dysfunction: research, mechanisms, and clinical implications. *J Man Manip Ther*. 2013;21(2):90–102. PubMed ID: [24421619](#) doi:[10.1179/2042618612Y.0000000027](#)
5. Goshtigian GR, Swanson BT. Using the selective functional movement assessment and regional interdependence theory to guide treatment of an athlete with back pain: a case report. *Int J Sports Phys Ther*. 2016;11(4):575–595. PubMed ID: [27525182](#)
6. Mokha GM, Sprague PA, Rodriguez R, Gatens DR. Functional movement pattern training improves mechanics in a female runner with external snapping hip syndrome. *Int J Athl Ther Train*. 2015; 20(1):25–33. doi:[10.1123/ijatt.2014-0095](#)
7. Busch AM, Clifton DR, Onate JA, Ramsey VK, Cromartie F. Relationship of preseason movement screens with overuse symptoms in collegiate baseball players. *Int J Sports Phys Ther*. 2017;12(6): 960–966. PubMed ID: [29158957](#) doi:[10.26603/ijst20170960](#)
8. Davenport KL, Air M, Grierson MJ, Krabak BJ. Examination of static and dynamic core strength and rates of reported dance related injury in collegiate dancers: a cross-sectional study. *J Dance Med Sci*. 2016; 20(4):151–161. doi:[10.12678/1089-313X.20.4.151](#)
9. Allen N, Nevill AM, Brooks JHM, Koutedakis Y, Wyon MA. The effect of a comprehensive injury audit program on injury incidence in ballet: a 3-year prospective study. *Clin J Sport Med*. 2013;23(5): 373–378. PubMed ID: [23558333](#) doi:[10.1097/JSM.0b013e3182887f32](#)
10. Krzyzanowicz R, Baker R, Nasypany A, Gargano F, Seegmiller J. Patient outcomes utilizing the selective functional movement assessment and mulligan mobilizations with movement on recreational dancers with sacroiliac joint pain: a case series. *Int J Athl Ther Train*. 2015;20(3):31–37. doi:[10.1123/ijatt.2014-0066](#)
11. Glaws KR, Juneau CM, Becker LC, Di Stasi SL, Hewett TE. Intra- and inter-rater reliability of the selective functional movement assessment (SFMA). *Int J Sports Phys Ther*. 2014;9(2):195–207. PubMed ID: [24790781](#)
12. Oxford Centre of Evidence-Based Medicine. CEBM levels of evidence 2.1. 2011. <http://www.cebm.net/wp-content/uploads/2014/06/CEBM-Levels-of-Evidence-2.1.pdf>. Accessed December 14, 2017.