

Should the Equilibrium Point Hypothesis (EPH) be Considered a Scientific Theory?

Robert L. Sainburg
Penn State University

The purpose of this commentary is to discuss factors that limit consideration of the equilibrium point hypothesis as a scientific theory. The EPH describes control of motor neuron threshold through the variable λ , which corresponds to a unique referent configuration for a muscle, joint, or combination of joints. One of the most compelling features of the equilibrium point hypothesis is the integration of posture and movement control into a single mechanism. While the essential core of the hypothesis is based upon spinal circuitry interacting with peripheral mechanics, the proponents have extended the theory to include the higher-level processes that generate λ , and in doing so, imposed an injunction against the supraspinal nervous system modeling, computing, or predicting dynamics. This limitation contradicts evidence that humans take account of body and environmental dynamics in motor selection, motor control, and motor adaptation processes. A number of unresolved limitations to the EPH have been debated in the literature for many years, including whether muscle resistance to displacement, measured during movement, is adequate to support this form of control, violations in equifinality predictions, spinal circuits that alter the proposed invariant characteristic for muscles, and limitations in the description of how the complexity of spinal circuitry might be integrated to yield a unique and stable equilibrium position for a given motor neuron threshold. In addition, an important empirical limitation of EPH is the measurement of the invariant characteristic, which needs to be done under a constant central state. While there is no question that the EPH is an elegant and generative hypothesis for motor control research, the claim that this hypothesis has reached the status of a scientific theory is premature.

Keywords: motor control, neural control of movement

In the target article, the author states, “There is only one coherent physical (physiological) theory that tries to identify the laws of physics that account for the neural control of biological movement.” While the equilibrium point hypothesis has played an important role in the science of motor control, it remains largely speculative, and cannot be considered a “scientific theory” because of substantial limitations in both its scope and because of controversial and largely unsupported features of the hypothesis. The definition of scientific theory published by AAAS

The author is with the Dept. of Kinesiology, Penn State University, University Park, PA. Address author correspondence to Robert L. Sainburg at rls45@psu.edu.

is “a well-substantiated explanation of some aspect of the natural world that can incorporate facts, laws, inferences, and tested hypotheses... In science, theories do not turn into facts through the accumulation of evidence... [Theories] incorporate a large body of scientific facts, laws, tested hypotheses, and logical inferences” (2007). It is important to emphasize that scientific theories are inductive in nature and aim for predictive and explanatory force. Therefore, I consider the author’s premise overly ambitious for the EP hypothesis of motor control. The purpose of the following discussion is not to critique the validity of the Equilibrium Point Hypothesis (EPH), but rather to discuss some limitations to its consideration as a scientific theory, particularly in its predictive and explanatory power.

While the EPH may, at first glance, seem appropriate to the description of scientific theory given above, there are many factors that prevent EPH from realizing the required predictive and explanatory power. EPH is an elegant hypothesis that combines the physical behavior of the musculoskeletal system with selected but limited elements of spinal anatomy and physiology. Those elements focus heavily on the reflex circuits associated with the muscle spindle, specifically those involving the intrafusal fibers that transduce small changes in muscle length. Together with the short and long loop components of the stretch reflex, this feedback has the potential to influence muscle activation, depending on the state of the specific motor neurons that receive input, either directly or indirectly from muscle afferents. According to the EPH, motor neuron threshold is specified through descending commands, and corresponds to a unique muscle length, for a given mechanical condition in the environment. This variable is referred to as λ . Because of the hypothesized invariant relationship between muscle force and muscle length across the full physiological range of the muscle, a unique referent configuration can be specified by λ , while the actual configuration depends also on external forces. Therefore, neural control of movement can be simplified to specification of this threshold-determining variable. One of the most compelling features of the equilibrium point hypothesis is the integration of posture and movement control into a single mechanism. Thus, at the end of motion and in the absence of external forces, a new posture is achieved without persistent muscle activity. Such behavior is best explained by a mechanism which modifies a position-dependent threshold, such as proposed by the EPH.

The EPH was introduced in a seminal series of papers by Feldman and colleagues (Asatrian & Fel’dman, 1965, Fel’dman, 1966a,b), which demonstrated that the force developed by a muscle, when the subject is instructed to maintain a particular angle, is a function of the length of the muscle and the variable λ . λ reflected an empirically derived value, the minimum length at which the passively lengthened muscle begins to be excited when the subject is asked not to intervene. These papers introduced the concept of an invariant characteristic for the muscle, or the force/length relationship for a given instructed state, presumably reflecting the centrally specified value, λ . Further, these early papers generalized the concept from a muscle to a group of muscles acting about a joint. Later studies generalized the model to multiple joints (Flash, 1987), and whole body postures as well as to locomotor control (Feldman et al., 2011).

From a neuroanatomical perspective, the equilibrium point hypothesis has two essential levels of consideration: spinal circuitry that provides the basis for λ as a control variable, and supraspinal processes that generate λ . While the

essential core of the hypothesis is based upon spinal circuitry interacting with peripheral mechanics, the proponents, largely by inference, have extended the theory to include the higher-level processes that generate descending commands to the spinal cord (Ostry & Feldman, 2003). In fact, an essential limitation to EPH is the injunction imposed by its proponents against the supraspinal nervous system modeling, computing, or predicting dynamics. This point seems to reflect a philosophical preference of the authors, rather than a rigorous consideration of the EP model. There is no question that calculations of formal dynamics using symbolic math do not occur in the CNS, except through declarative cognitive processes. Nevertheless, the evidence that humans take account of dynamic factors for motor selection and motor control is overwhelming. This includes evidence from adaptation studies showing generalization of dynamics and aftereffects reflecting previously applied loads (Dizio & Lackner, 1995, Conditt et al., 1997, Ghez et al., 1999, Sainburg et al., 1999, Hinder & Milner, 2003, Hwang & Shadmehr, 2005), evidence that humans often select actions that have reduced energetic requirements (Alexander, 1997, Massaad et al., 2007, Nishii & Tani, 2009), evidence for hemispheric asymmetries in accounting for dynamics in coordination of reaching, an effect that is hard to explain if the central nervous system does not account for dynamics (Sainburg & Kalakanis, 2000, Bagesteiro & Sainburg, 2002, Sainburg 2002, Sainburg & Schaefer, 2004, Schabowsky et al., 2007, Schaefer et al., 2007, Schaefer et al., 2009, Mutha et al., 2012, Przybyla et al., 2012, Schaefer et al., 2012, Coelho et al., 2013, Mani et al., 2013), evidence that we account for the effects of inertial dynamics when making point to point targeted movements (Ghez et al., 1990, Sainburg et al., 1993, Gordon et al., 1994a,b, Ghez & Sainburg 1995, Sainburg et al., 1995), and evidence that we predict object behavior based on physical properties (Dingwell et al., 2002, Dingwell et al., 2004, Cothros et al., 2006, Kluzik et al., 2008, Chib et al., 2009). None of these observations precludes the idea that movement must ultimately be controlled by specification of joint, limb, and body referent configurations. However, the EPH proponents find the idea of internal models of physical phenomenon to be distasteful, seemingly due to philosophical reasons.

There are a variety of unresolved limitations to this hypothesis that have been debated in the literature for many years, including whether muscle resistance to displacement (often referred to as muscle stiffness) measured during movement is large enough to support this form of control (Gomi & Kawato, 1997, Popescu & Rymer, 2000, Hinder & Milner, 2003, Popescu & Rymer, 2003), and violations in equifinality predictions (Popescu & Rymer, 2000, Sainburg et al., 2003). However, it should be stressed that the assumptions of equifinality are based on nonbiological mass-spring systems, and should not be assumed for all cases of biological control because muscle force-length properties, as well as sensorimotor responses, are largely nonlinear (see Feldman & Latash, 2005). Nichols and colleagues have recently demonstrated that heterogeneous inhibitory force feedback from tendon organs can alter the slope of the muscle length tension relationship, calling into question its invariance and relegating a single threshold/length variable (λ) inadequate to support control (Nichols & Ross, 2009). With regard to the complexity and variety of spinal circuits (Tresch et al., 1999, Loeb, 2001), the EPH has no explicit description of how all these might be integrated to yield a unique and stable equilibrium position for a given motor neuron threshold.

An important empirical limitation of EPH is the measurement of the invariant characteristic, which needs to be done under a constant central state. However, there has been no formal method to assure this can be done. In humans, the strategy has been to tell people, ‘do not intervene’ (or some similar instruction), and assume that this stabilizes central commands to muscles (Latash, 1992, Archambault et al., 2005). A common, though anecdotal, demonstration of referent configuration involves the unloading response: a weight, such as a book supported on the hand with the forearm held out straight, is removed from the hand. The typical response is that the elbow immediately flexes and achieves a new, more flexed, position. This position is described as the specified referent configuration for the elbow when holding the book. However, an alternative explanation is that the static equilibrium between elbow flexor force, and gravitational load is disrupted, producing elbow flexor acceleration. Flexor motion elicits an extensor stretch reflex, which brings the limb to a stop and allows the subject to achieve a new equilibrium between elbow force and gravitational load. This sequence does not require a predetermined referent configuration. The argument against this interpretation is that subjects were told not to intervene, which may or may not be effective in preventing voluntary actions. However, the effects described do not require declarative cognitive mediation. In fact, a demonstration that argues against the EPH description of the unloading experiment is a condition in which a very large load is held by near maximal elbow torque, such as a heavy dumbbell. The required referent configuration for such a large torque would be outside of the anatomical range of the limb. Nevertheless, unloading the arm does not accelerate the limb into its full range, even if people are told not to intervene. Rather a new position is achieved in an intermediate range. Therefore, even though subjects are told, “do not intervene”, their limb is not displaced to the extreme of the anatomical range. Instead, subjects achieve a comfortable posture, which certainly does not reflect the referent configuration for supporting the dumbbell before unloading.

In summary, the EPH has proved to be an elegant and generative hypothesis in motor control research. It is currently the only hypothesis that resolves the posture-movement paradox, integrating the control of posture and movement into a single control scheme. However, the claim that this hypothesis has reached the status of a scientific theory seems premature. Many aspects of motor control remain unexplained by EPH, and some phenomena appear to contradict this hypothesis.,. Therefore, the predictive and explanatory power of EPH remains too limited to consider it an accepted scientific theory, at this point in time.

References

- Alexander, R.M. (1997). A minimum energy cost hypothesis for human arm trajectories. *Biological Cybernetics*, 76(2), 97–105. PubMed doi:10.1007/s004220050324
- AAAS (2007). The Role of Theory in Advancing 21st Century Biology: Catalyzing Transformative Research.
- Archambault, P.S., Mihaltchev, P., Levin, M.F., & Feldman, A.G. (2005). Basic elements of arm postural control analyzed by unloading. *Experimental Brain Research*, 164(2), 225–241. PubMed doi:10.1007/s00221-005-2245-6
- Asatryan, D.G., & Fel’dman, A.G. (1965). [On the functional structure of the nervous system during movement control or preservation of a stationary posture. I. Mechanographic

- analysis of the action of a joint during the performance of a postural task]. *Biofizika*, 10(5), 837–846. [PubMed](#)
- Bagesteiro, L.B., & Sainburg, R.L. (2002). Handedness: dominant arm advantages in control of limb dynamics. *Journal of Neurophysiology*, 88(5), 2408–2421. [PubMed doi:10.1152/jn.00901.2001](#)
- Chib, V.S., Krutky, M.A., Lynch, K.M., & Mussa-Ivaldi, F.A. (2009). The separate neural control of hand movements and contact forces. *The Journal of Neuroscience*, 29(12), 3939–3947. [PubMed doi:10.1523/JNEUROSCI.5856-08.2009](#)
- Coelho, C.J., Przybyla, A., Yadav, V., & Sainburg, R.L. (2013). Hemispheric differences in the control of limb dynamics: a link between arm performance asymmetries and arm selection patterns. *Journal of Neurophysiology*, 109(3), 825–838. [PubMed doi:10.1152/jn.00885.2012](#)
- Conditt, M.A., Gandolfo, F., & Mussa-Ivaldi, F.A. (1997). The motor system does not learn the dynamics of the arm by rote memorization of past experience. *Journal of Neurophysiology*, 78(1), 554–560. [PubMed](#)
- Cothros, N., Wong, J.D., & Gribble, P.L. (2006). Are there distinct neural representations of object and limb dynamics? *Experimental Brain Research*, 173(4), 689–697. [PubMed doi:10.1007/s00221-006-0411-0](#)
- Dingwell, J.B., Mah, C.D., & Mussa-Ivaldi, F.A. (2002). Manipulating objects with internal degrees of freedom: evidence for model-based control. *Journal of Neurophysiology*, 88(1), 222–235. [PubMed](#)
- Dingwell, J.B., Mah, C.D., & Mussa-Ivaldi, F.A. (2004). Experimentally confirmed mathematical model for human control of a non-rigid object. *Journal of Neurophysiology*, 91(3), 1158–1170. [PubMed doi:10.1152/jn.00704.2003](#)
- Dizio, P., & Lackner, J.R. (1995). Motor adaptation to Coriolis force perturbations of reaching movements: endpoint but not trajectory adaptation transfers to the nonexposed arm. *Journal of Neurophysiology*, 74(4), 1787–1792. [PubMed](#)
- Fel'dman, A.G. (1966a). [On functional tuning of nervous system during controlled or preservation of stationary pose. 3. Mechanographic analysis of human performance of simple movement tasks]. *Biofizika*, 11(4), 667–675. [PubMed](#)
- Fel'dman, A.G. (1966b). [On the functional tuning of the nervous system in movement control or preservation of stationary pose. II. Adjustable parameters in muscles]. *Biofizika*, 11(3), 498–508. [PubMed](#)
- Feldman, A.G., Krasovskiy, T., Banina, M.C., Lamontagne, A., & Levin, M.F. (2011). Changes in the referent body location and configuration may underlie human gait, as confirmed by findings of multi-muscle activity minimizations and phase resetting. *Experimental Brain Research*, 210(1), 91–115. [PubMed doi:10.1007/s00221-011-2608-0](#)
- Flash, T. (1987). The control of hand equilibrium trajectories in multi-joint arm movements. *Biological Cybernetics*, 57(4-5), 257–274. [PubMed doi:10.1007/BF00338819](#)
- Ghez, C., Gordon, J., Ghilardi, M.F., Christakos, C.N., & Cooper, S.E. (1990). Roles of proprioceptive input in the programming of arm trajectories. *Cold Spring Harbor Symposia on Quantitative Biology*, 55, 837–847. [PubMed doi:10.1101/SQB.1990.055.01.079](#)
- Ghez, C., Krakauer, J., Sainburg, R.L., & Ghilardi, M.F. (1999). *Spatial representations and internal models of limb dynamics in motor learning*. *The New Cognitive Sciences*. M. S. Gazzaniga. Cambridge, MA: The MIT Press.
- Ghez, C., & Sainburg, R. (1995). Proprioceptive control of interjoint coordination. *Canadian Journal of Physiology and Pharmacology*, 73(2), 273–284. [PubMed doi:10.1139/y95-038](#)
- Gomi, H., & Kawato, M. (1997). Human arm stiffness and equilibrium-point trajectory during multi-joint movement. *Biological Cybernetics*, 76(3), 163–171. [PubMed doi:10.1007/s004220050329](#)
- Gordon, J., Ghilardi, M.F., Cooper, S.E., & Ghez, C. (1994a). Accuracy of planar reaching movements. II. Systematic extent errors resulting from inertial anisotropy. *Experimental Brain Research*, 99(1), 112–130. [PubMed doi:10.1007/BF00241416](#)

- Gordon, J., Ghilardi, M.F., & Ghez, C. (1994b). Accuracy of planar reaching movements. I. Independence of direction and extent variability. *Experimental Brain Research*, 99(1), 97–111. [PubMed doi:10.1007/BF00241415](#)
- Hinder, M.R., & Milner, T.E. (2003). The case for an internal dynamics model versus equilibrium point control in human movement. *The Journal of Physiology*, 549(Pt 3), 953–963. [PubMed doi:10.1113/jphysiol.2002.033845](#)
- Hwang, E.J., & Shadmehr, R. (2005). Internal models of limb dynamics and the encoding of limb state. *Journal of Neural Engineering*, 2(3), S266–S278. [PubMed doi:10.1088/1741-2560/2/3/S09](#)
- Kluzik, J., Diedrichsen, J., Shadmehr, R., & Bastian, A.J. (2008). Reach adaptation: what determines whether we learn an internal model of the tool or adapt the model of our arm? *Journal of Neurophysiology*, 100(3), 1455–1464. [PubMed doi:10.1152/jn.90334.2008](#)
- Latash, M.L. (1992). Independent control of joint stiffness in the framework of the equilibrium-point hypothesis. *Biological Cybernetics*, 67(4), 377–384. [PubMed doi:10.1007/BF02414893](#)
- Loeb, G.E. (2001). Learning from the spinal cord. *The Journal of Physiology*, 533(Pt 1), 111–117. [PubMed doi:10.1111/j.1469-7793.2001.0111b.x](#)
- Mani, S., Mutha, P.K., Przybyla, A., Haaland, K.Y., Good, D.C., & Sainburg, R.L. (2013). Contralateral motor deficits after unilateral stroke reflect hemisphere-specific control mechanisms. *Brain*, 136(Pt 4), 1288–1303. [PubMed doi:10.1093/brain/aws283](#)
- Massaad, F., Lejeune, T.M., & Detrembleur, C. (2007). The up and down bobbing of human walking: a compromise between muscle work and efficiency. *The Journal of Physiology*, 582(Pt 2), 789–799. [PubMed doi:10.1113/jphysiol.2007.127969](#)
- Mutha, P.K., Haaland, K.Y., & Sainburg, R.L. (2012). The effects of brain lateralization on motor control and adaptation. *Journal of Motor Behavior*, 44(6), 455–469. [PubMed doi:10.1080/00222895.2012.747482](#)
- Nichols, R., & Ross, K.T. (2009). The implications of force feedback for the lambda model. *Advances in Experimental Medicine and Biology*, 629, 663–679. [PubMed doi:10.1007/978-0-387-77064-2_36](#)
- Nishii, J., & Tani, Y. (2009). Evaluation of trajectory planning models for arm-reaching movements based on energy cost. *Neural Computation*, 21(9), 2634–2647. [PubMed doi:10.1162/neco.2009.06-08-798](#)
- Ostry, D.J., & Feldman, A.G. (2003). A critical evaluation of the force control hypothesis in motor control. *Experimental Brain Research*, 153(3), 275–288. [PubMed doi:10.1007/s00221-003-1624-0](#)
- Popescu, F.C., & Rymer, W.Z. (2000). End points of planar reaching movements are disrupted by small force pulses: an evaluation of the hypothesis of equifinality. *Journal of Neurophysiology*, 84(5), 2670–2679. [PubMed](#)
- Popescu, F.C., & Rymer, W.Z. (2003). Implications of low mechanical impedance in upper limb reaching motion. *Motor Control*, 7(4), 323–327. [PubMed](#)
- Przybyla, A., Good, D.C., & Sainburg, R.L. (2012). Dynamic dominance varies with handedness: reduced interlimb asymmetries in left-handers. *Experimental Brain Research*, 216(3), 419–431. [PubMed doi:10.1007/s00221-011-2946-y](#)
- Sainburg, R.L. (2002). Evidence for a dynamic-dominance hypothesis of handedness. *Experimental Brain Research*, 142(2), 241–258. [PubMed doi:10.1007/s00221-001-0913-8](#)
- Sainburg, R.L., Ghez, C., & Kalakanis, D. (1999). Intersegmental dynamics are controlled by sequential anticipatory, error correction, and postural mechanisms. *Journal of Neurophysiology*, 81(3), 1045–1056. [PubMed](#)
- Sainburg, R.L., Ghilardi, M.F., Poizner, H., & Ghez, C. (1995). Control of limb dynamics in normal subjects and patients without proprioception. *Journal of Neurophysiology*, 73(2), 820–835. [PubMed](#)

- Sainburg, R.L., & Kalakanis, D. (2000). Differences in control of limb dynamics during dominant and nondominant arm reaching. *Journal of Neurophysiology*, *83*(5), 2661–2675. [PubMed](#)
- Sainburg, R.L., Lateiner, J.E., Latash, M.L., & Bagesteiro, L.B. (2003). Effects of altering initial position on movement direction and extent. *Journal of Neurophysiology*, *89*(1), 401–415. [PubMed](#) doi:10.1152/jn.00243.2002
- Sainburg, R.L., Poizner, H., & Ghez, C. (1993). Loss of proprioception produces deficits in interjoint coordination. *Journal of Neurophysiology*, *70*(5), 2136–2147. [PubMed](#)
- Sainburg, R.L., & Schaefer, S.Y. (2004). Interlimb differences in control of movement extent. *Journal of Neurophysiology*, *92*(3), 1374–1383. [PubMed](#) doi:10.1152/jn.00181.2004
- Schabowsky, C.N., Hidler, J.M., & Lum, P.S. (2007). Greater reliance on impedance control in the nondominant arm compared with the dominant arm when adapting to a novel dynamic environment. *Experimental Brain Research*, *182*(4), 567–577. [PubMed](#) doi:10.1007/s00221-007-1017-x
- Schaefer, S.Y., Haaland, K.Y., & Sainburg, R.L. (2007). Ipsilesional motor deficits following stroke reflect hemispheric specializations for movement control. *Brain*, *130*(Pt 8), 2146–2158. [PubMed](#) doi:10.1093/brain/awm145
- Schaefer, S.Y., Haaland, K.Y., & Sainburg, R.L. (2009). Hemispheric specialization and functional impact of ipsilesional deficits in movement coordination and accuracy. *Neuropsychologia*, *47*(13), 2953–2966. [PubMed](#) doi:10.1016/j.neuropsychologia.2009.06.025
- Schaefer, S.Y., Mutha, P.K., Haaland, K.Y., & Sainburg, R.L. (2012). Hemispheric specialization for movement control produces dissociable differences in online corrections after stroke. *Cerebral Cortex*, *22*(6), 1407–1419. [PubMed](#) doi:10.1093/cercor/bhr237
- Tresch, M.C., Saltiel, P., & Bizzi, E. (1999). The construction of movement by the spinal cord. *Nature Neuroscience*, *2*(2), 162–167. [PubMed](#) doi:10.1038/5721